

$\beta - P$ Connectedness in L -Bitopological spaces

FangLi Li , Hongkui Li, Tao Li

Abstract— This paper introduce open sets and p - closed sets into L -Bitopological spaces, and based on this we introduce some related definitions and theorems about $\beta - P$ closed set and $\beta - P$ open sets . Furthermore, we give a new concept about $\beta - P$ local-connectivity .Then, it point out $\beta - P$ local-connectivity has two properties of topological invariance and finitely productive property and proves the other relevant theories.

Index Terms— L - bitopological spaces, $\beta - P$ local-connectivity, topological invariance, finitely productive property

I. INTRODUCTION

Since J. C. Kelly introduced the concept of double topological space, many scholars has a great interest about researching dual topology, then introduced L -bitopological spaces on the basis of L -topological space, and makes a long-term study of the separation .Connectivity is an important branch of fuzzy topology, the domestic scholars have studied the variety of connectivity, such as θ -connectivity[3], connectivity[4], stratified connectedness [5], disconnectedness branches e .c. Due to the concept of connectivity is closely related to geometric closure , many connectivity is also defined based on the definition of different closure concepts. In this paper,we introduced $\beta - P$ open sets and $\beta - P$ closed sets in L -Bitopological spaces on the basis of $\beta - P$ open sets and $\beta - P$ closed sets .In this case ,we defined a new connectivity in L - bi-topological spaces which is called $\beta - P$ local-connectivity ,and the study of the connectivity has gotten some good properties.

II. PRELIMINARY KNOWLEDGE

A. L -bitopological spaces

Definition 2.1: Let L be a F lattice ,that is a completely distributive lattice with the reverse involution ,let X be a common set and let L^X is a set that contains the whole L - fuzzy sets on X ,0 and 1 respectively expressed the minimum and maximum in L , $M(L)$ and $M^*(L^X)$ respectively expressed all molecules of L and L^X ,we record

L -bts as L - bitopological space and $A_{\delta_1^-}$ as the closure of A in (L^X, δ_1) [6].

B. Final Stage The Related Defines And Conclusion About p - open set And p - close Set

Definition 2.2: Let (L^X, δ) be L -bts , $A \in L^X$ and record as p - open set ,if and only if have open set U ,makes $A \leq U \leq A^-$;If A is a p - open set, we call A' is a p - close set .The whole p - open sets in (L^X, δ) are denoted by $LPO(L^{X*})$,and the whole p - close sets are denoted by $LPC(L^{X*})$ [7].

Note: The close set in L -bts must be p - close set, and on the contrary generally does not set up.

Lemma 2.1: Let (L^X, δ) be L -bts , then $\delta \subset LPO(L^{X*})$, $\delta' \subset LPC(L^{X*})$.

Definition 2.3: Let (L^X, δ) be L -bts , $A \in L^{X*}$, $B \in L^{X*}$, then:

(a) The union of all p - open sets which contained by A is called internal of A 's $LE - p^*$,record as $A^{*\Delta}$,that is

$$A^{*\Delta} = \{B \in LPO(L^{X*}) | B \leq A\}.$$

(b) The union of all p - close sets which contained by A is called external of A 's $LE - p^*$,record as $A^{*\leftarrow}$,that is

$$A^{*\leftarrow} = \{B \in LPC(L^{X*}) | A \leq B\}$$

Lemma 2.2: Let (L^X, δ) be L -bts , $A, B \in L^{X*}$,then

(a) $A^\circ \leq A^{*\Delta} \leq A \leq A^{*\leftarrow} \leq A^-$.

(b) If $A \in \delta \cap LPC(L^{X*})$,then $A \in \delta'$.

(c) If $A \in \delta \cap LPO(L^{X*})$,then $A \in \delta$.

(d) If $A \leq B$,then $A^{*\leftarrow} \leq B^{*\leftarrow}$, $A^{*\Delta} \leq B^{*\Delta}$.

(e) $(A \vee B)^{*\leftarrow} = A^{*\leftarrow} \vee B^{*\leftarrow}$, $(A \vee B)^{*\Delta} = A^{*\Delta} \vee B^{*\Delta}$.

(f) The arbitrary intersection of p - closed sets is p - closed set, and the arbitrary intersection of p - open sets is p - open set.

FangLi Li, School of Science, Shandong University of Technology, Zibo, China

Hongkui Li, School of Science, Shandong University of Technology, Zibo, China.

Tao Li, Innovation and entrepreneurship College, Shandong University of Technology, Zibo, China.

C. Figures The Introduction of $\beta - p$ open set And $\beta - p$ close set

According to the preparation, we introduce $\beta - p$ open set and $\beta - p$ close set into L — bitopological spaces, and give some related properties and theorems [7] about $\beta - p$ open set and $\beta - p$ close set on the basis of p — closed set and p — open set, and give the corresponding proof.

Definition 3.1: Let (L^X, δ) be L — bts , $\beta \in L$, $A \in L^X$, record as $l_\alpha(A) = \{x \in X | A(x) > \beta\}$, let $\beta \in L - \{0\}$, if $l_\beta(A^{*\leftarrow}) = l_\beta(A)$, then call A as $\beta - p$ open set; If A' is $\beta - p$ close set, then we call A as $\beta - p$ open set.

Lemma 3.1: Let (L^X, δ) be L — bts , $A \in L^X$, if A is $\beta - p$ close set, then A is $\beta - p$ open set.

Proof: If A is $\beta - p$ close set, then about $\beta \in L - \{0\}$, $A \in L^X$, when $\gamma \geq \beta$, there is $l_\beta(A^-) = l_\beta(A)$, due to $A^{*\leftarrow} \leq A^{*\leftarrow}$, then there is $l_\beta(A^{*\leftarrow}) = l_\beta(A)$, so A is a $\beta - p$ open set.

Lemma 3.2: Let (L^X, δ) be L — bts , $A \in L^X$, if A is a $\beta - p$ open set, then A is $\beta - p$ open set.

Lemma 3.3: Let (L^X, δ) be L — bts , $A \in L^X$, if A is a p — close set is the necessary and sufficient condition of A is $1 - p$ close set.

Proof :

" \Leftarrow " Clearly established.

" \Rightarrow " Due to " A " is $1 - p$ close set, so $\forall \beta \geq 1' = 0, l_\beta(A^{*\leftarrow}) = l_\beta(A)$

If $A^{*\leftarrow} \neq A$, then there is $x \in X$, cause $A^{*\leftarrow}(x) > A(x)$, marked $\beta_0 = A(x) \geq 0$, then $x \notin l_{\beta_0}(A^{*\leftarrow})$ is conflict with $(*)$, so $A^{*\leftarrow} = A$, that is A is $\beta - p$ close set.

Lemma 3.4: Let (L^X, δ) be L — bts , $A, B \in L^X$, $\beta \in L - \{0\}$, then

- (a) If $A \leq B$, then $l_\alpha(A) \subseteq l_\alpha(B)$.
- (b) $l_\alpha(A \wedge B) = l_\alpha(A) \cap l_\alpha(B)$.
- (c) $l_\alpha((A \wedge B)^{*\Delta}) = l_\alpha(A^{*\Delta}) \cap l_\alpha(B^{*\Delta})$.
- (d) $l_\alpha((A \vee B)^{*\Delta}) = l_\alpha(A^{*\Delta}) \cup l_\alpha(B^{*\Delta})$.
- (e) $l_\alpha(A^{*\leftarrow} \wedge B^{*\leftarrow}) = l_\alpha(A^{*\leftarrow}) \cap l_\alpha(B^{*\leftarrow})$.

$$(f) l_\alpha(A^{*\leftarrow} \vee B^{*\leftarrow}) = l_\alpha(A^{*\leftarrow}) \cup l_\alpha(B^{*\leftarrow}).$$

D. $\beta - p$ local-connectivity

1. Definitions about $\beta - p$ local-connectivity

Definition 4.1.1: Let (L^X, δ) be L — bts , $A, B \in L^X$, $\alpha \in L - \{0\}$, if $A^{*\leftarrow} \wedge B \leq \beta'$ and $A \wedge B^{*\leftarrow} \leq \beta'$, then call A and B are $\beta - p$ insular.

Definition 4.1.2: Let (L^X, δ) be L — bts , $S \in L^X$, if there is no $A, B \in L^X$, to make A and B are $\beta - p$ insular, and $A \vee B = S$, $A > \beta'$, $B > \beta'$, then call S is Connected set in (L^X, δ) . Particularly, when 1_x which is the Maximum element in L^X is $\beta - p$ Connected set; Otherwise, call (L^X, δ) is $\beta - p$ disconnected space.

Definition 4.1.3: Let (L^X, δ) be L — bts , $x \in L^X$, if every neighborhood of A contains a $\beta - p$ connected neighborhood V , then call x is $\beta - p$ local-connectivity; Otherwise, call x is not $\beta - p$ local-connectivity [8].

Definition 4.1.4: If every point of (L^X, δ) is $\beta - p$ local-connectivity, then call (L^X, δ) is $\beta - p$ local-connected space.

2. Basic properties of $\beta - p$ local-connectivity

Theorem 4.2.1: The local connected space must be $\beta - p$ local-connected space.

Proof : From the definition of local-connected spaces, we can know that every neighborhood of A contains a $\beta - p$ connected neighborhood V , the V is open set. By definition 4.2 in the literature [3], we can know that the connected open set must be $\beta - p$ open set of V . As for $\forall x \in L^X$, if every neighborhood of A contains a $\beta - p$ connected neighborhood V , then (L^X, δ) is $\beta - p$ local-connected space according to definition 4.1.4.

On the contrary, does not necessarily set up.

Theorem 4.2.2: Let (L^X, δ) be L — bts , then the following conditions are equivalent:

- (a) L^X is a $\beta - p$ local-connected space.
- (b) Arbitrary $\beta - p$ connected branch of L^X 's arbitrary $\beta - p$ open set is a $\beta - p$ open set.
- (c) L^X is a connected base.

Proof :

(a) \Rightarrow (b):

Let U is a arbitrary $\beta - p$ open set of $\beta - p$ connected space, c is arbitrary $\beta - p$ connected branch of U , as for arbitrary $x \in c \in U$, then $U_x \in U$. And because

L^X is $\beta-p$ local-connected, there is a connected neighborhood V , and V also is $\beta-p$ connected neighborhood of subspace U [9].

\Rightarrow (c) :

Let α is a set family that formed by all $\beta-p$ open set of L^X , then α is a base of L^X , also because $\beta-p$ connected branch is $\beta-p$ connected subset, then α is connected base of L^X .

\Rightarrow (a) :

Let L^X has a $\beta-p$ connected base α , then each member of α are all $\beta-p$ connected sets, as for $\forall x \in L^X$, make $\alpha_x = \{A \in \beta | x \in A\}$, then $x \in \alpha_x \subset \alpha$, that α_x is $\beta-p$ connected, we can know L^X is $\beta-p$ local-connected by the definition 4.1.3.

Theorem 4.2.3: Let (L^X, δ_1) is a $\beta-p$ local-connected space, (X, T) is a topological space, $f: L^X \rightarrow X$ is a continuous open mapping, then we can construct that (L^X, δ_2) is a $\beta-p$ local-connected space by (X, T) .

Proof : Let L^X is a $\beta-p$ local-connected space, then we can know there is a $\beta-p$ connected base in L^X by the definition 4.2.2. As for f is a continuous open mapping, to make $\beta_0 = \{f(B) | B \in \beta\}$, then $\forall B \in \beta, f(B), \forall B \in \beta, f(B)$ is $\beta-p$ connected open set of X , so β_0 is $\beta-p$ connected open set family of X . As for arbitrary $\beta-p$ open set of X , f is $\beta-p$ connected and surjection, so:

$$A = f(f^{-1}(A)) = f\left(\bigcup_{B \in \beta_1} B\right) = \bigcup_{B \in \beta_1} f(B)$$

Then β_0 is a $\beta-p$ open connected base of X .

By the theorem 4.2.2, we can know (L^X, δ_2) can structured by (X, T) is $\beta-p$ connected space.

We call the nature is topological invariance which is can keep the same nature under $\beta-p$ continuous mapping in topological space.

Theorem 4.2.4: If $L^{X_1}, L^{X_2}, \dots, L^{X_n}$ are also $\beta-p$ local-connected space, then $L^X = L^{X_1} \times L^{X_2} \times \dots \times L^{X_n}$ is $\beta-p$ local-connected space too.

Proof : Let $L^{X_i} (i=1, 2, \dots, n)$ is $\beta-p$ connected space, from theorem 4.2.2, we can know that there is a $\beta-p$ connected base $V_i (i=1, 2, \dots, n)$ in L^{X_i} , then make

$V = \{V_1 \times V_2 \times \dots \times V_n | V_i \in V (i=1, 2, \dots, n)\}$, and from literature [10], we can know $\beta-p$ connected nature is finitely productive property, so V is a $\beta-p$ connected base of product space L^X . Also, from theorem 4.2.2, we can know that product space $L^X = L^{X_1} \times L^{X_2} \times \dots \times L^{X_n}$ is also a $\beta-p$ local-connected space.

Some properties P of topological spaces are called finite integrable properties, if there are $n \geq 1$ arbitrary topological space $L^{X_1}, L^{X_2}, \dots, L^{X_n}$ have property P , and contained product space $L^X = L^{X_1} \times L^{X_2} \times \dots \times L^{X_n}$ also have property P , then $\beta-p$ local-connected nature is finitely productive property.

REFERENCES

- [1] Kelly J C. Bitopological Spaces[J]. Proc London Math Soc, 1963(77): 71-89.
- [2] Zhang Chunbing, Zheng Chongyou. Separation axiom of L-double fuzzy topological space[J]. Journal of Beijing Teachers College (NATURAL SCIENCE EDITION), 1990.11(1): 13-17.
- [3] Liu Hongping, Meng Guangwu, Hu Xiaonan. The θ Connectivity of L -bitopological spaces. Journal of Shantou University (NATURAL SCIENCE EDITION), 2008.05.
- [4] Zhao Haixin, Meng Guangwu. The Connectivity of L -bitopological spaces[J]. Journal of Jinggangshan University (NATURAL SCIENCE EDITION), 2010.11.15.
- [5] Xu Hao, Meng Guangwu. The stratified connectedness of L -bitopological spaces[J]. Journal of Liaocheng University, 2010.06.25.
- [6] Wang Xiaoxia, Jiang Jinping. Several classes of weak-connectedness branches of L -bitopological spaces[D]. Journal of Yunnan Normal University, 2013.07.
- [7] Xu Xiaoling. The $\alpha-p$ Connectivity of L -fuzzy with L-double fuzzy topological space, 2009.06.
- [8] Xiong Jincheng. Point set topology notes[D]. Fourth edition, 2011.06.
- [9] Jiang Pingyuan, Wang Xiaoxia. θ -local connected space. Jiangxi Science. Volume 26, First period, 2008, 02.
- [10] Zhang Xiangting. Calculation of wind pressure and wind induced vibration of structures[M]. Shanghai: Tongji University press, 1985.

Corresponding Author: Hongkui Li

Acknowledgment: This work is supported by Doctoral Fund of Shandong University of Technology of China (4041/415061), Comprehensive budget management system design (9101/215043), Shandong Research project of teaching reform in undergraduate colleges and universities of China (2015Z071).